




藥物 常識

如何使用

Premarin® Vaginal Cream

How To Use

Premarin® Vaginal Cream




香港浸信會醫院
Hong Kong Baptist Hospital

如何使用 Premarin® Vaginal Cream

How To Use Premarin® Vaginal Cream

1. 清洗雙手。
Wash hands.
2. 把藥膏與投藥器接連，將投藥器之活塞拉出至醫生所指示的劑量刻度，然後按壓藥膏直至投藥器內的空間被完全注滿。
Connect the applicator with the tube. Pull out applicator plunger until the directed dosage mark. Press the ointment tube to fill up the applicator.
3. 仰臥床上，曲起雙膝，使投藥器較易插入。
Insertion of the applicator is more easily accomplished when lying on the back with knees bent.
4. 從藥膏解下投藥器，小心地把投藥器輕輕塞入陰道深處，按下活塞，藥膏便會留在陰道內。
Disconnect the applicator. Gently insert the applicator into the vagina as far as it will go comfortably. Depress the plunger to transfer the medicine to the vagina.



5. 使用後，將活塞從投藥器拉出，用溫水及肥皂清洗乾淨，弄乾並保持清潔。
After use, remove the plunger by pulling it out of the applicator. Then wash both parts in warm soapy water, rinse and dry carefully.



6. 此治療法最好在睡前進行。
This treatment is to be used preferably just before going to bed.
7. 病徵病狀可能會很快消失，但仍須按醫生指示繼續使用，直至整個療程完畢。
Itching, burning and discharge may disappear quickly but it is important that you continue the treatment until finished.

8. 此用法不適用於孕婦，如有疑問請聯絡醫生或藥房。

This method is not suitable for pregnant woman.

For enquiry, please contact your doctor or our pharmacy department.

此藥物指引只供參考之用。

用藥應按照醫生指示。

如有疑問請向主診醫生或藥房職員查詢。

藥物查詢熱線: 23398964

The above information provided can be regarded as a general reference. If you have any problem, please consult your doctor or your pharmacist.

Pharmacy hotline: 23398964

香港浸信會醫院網址: www.hkbh.org.hk

HK Baptist Hospital web site: www.hkbh.org.hk

香港浸信會醫院藥房部編制

Printed by Hong Kong Baptist Hospital Pharmacy Department