香港浸信會醫院
Hong Kong Baptist Hospital

肉毒桿菌 / 組織填充劑注射
Injection – Botulinum Toxin / Filler

醫療程序須知

簡介

肉毒桿菌 / 組織填充劑注射是非手術性的美型輪廓治療，效果非永久性的。

適應症

- 改善面部皺紋
- 嚼肌縮小
- 腋下及手部多汗症
- 塑造豐盈的面部
- 改善小腿線條
- 頸闊肌紋

手術 / 程序前須知

1. 可能需要局部麻醉。
2. 可能需要多次治療，而且未必能全部清除。
3. 改善程度並無任何保證，治療反應因人而異。
4. 治療均存在復發機會。
5. 治療前後均需拍照記錄，以作治療進度之參考。

手術 / 程序後須知

1. 注射部位會出現疼痛、瘀傷、腫脹乃屬常見現象。
2. 治療後可恢復使用護膚產品和化妝品。

風險及併發症

1. 一般風險：泛紅、腫脹、瘀傷、感染、過度矯正、矯正不足、不對稱效果、過敏反應。
2. 肉毒桿菌注射風險：感冒徵狀、眩暈、頭痛、肌肉無力、複視、眼皮下垂、眉弓下垂、眼乾。肉毒桿菌在多次重覆使用後效果會減弱。
3. 組織填充劑注射風險：皮膚結塊、皮膚壞死、失明。

免責聲明

本單張只提供有關手術 / 程序的基本資料，並列出常見的風險或併發症；可能發生的風險或併發症不能盡錄，亦可因人而異。如需詳細資料或個別查詢，請聯絡你的醫生。

參考資料

1. M Kassir. Complications of botulinum toxin and fillers: A narrative review. J Cosmet Derm 2020. 19(3):570-573
2. M Signorini et al. Global Aesthetics Consensus: Avoidance and Management of Complications from Hyaluronic Acid Fillers-Evidence- and Opinion-Based Review and Consensus Recommendations. Plast Reconstr Surg 2016.137(6):961e-971e
3. H Sundaram et al. Global Aesthetics Consensus: Botulinum Toxin Type A--Evidence-Based Review, Emerging Concepts, and Consensus Recommendations for Aesthetic Use, Including Updates on Complications. Plast Reconstr Surg 2016. 137(3):518e-529e

Patient's Label
Patient Name: _____
Hospital No:_____
Episode No:_____

病人簽署：_____ 簽署日期：_____