香港浸信會醫院
Hong Kong Baptist Hospital

聚焦超聲波
HIFU (Ulfit, Liftera V)

聚焦超聲波 HIFU (Ulfit, Liftera V)

醫療程序須知

簡介

「高能聚焦超聲波」利用高度集中的超聲波，將能量聚焦至淺層真皮層 / 深層真皮層 / 筋膜層，在減少傷及其他組織的情況下，令目標蛋白產生「變性作用」(Denaturation)，從而希望達致面頰提升、收緊雙下巴、收緊頸項、輪廓提升、減淡皺紋、緊緻肌膚、回復彈性等功效。當用於身體時，可能溶解並破壞脂肪細胞而改善局部多餘脂肪。

手術 / 程序前須知

1. 以下情況並不適合進行高能聚焦超聲波治療：治療部位有開放式傷口 / 創傷、嚴重或囊腫形暗瘡、皮膚發炎、增生性疤痕，孕婦，患有免疫系統疾病、癲癇症、治療部位有神經痲痺問題、凝血不正常、或皮膚癌病者。
2. 如有植入物、皮下填充劑、植入式電子儀器、金屬支架位於治療部位或附近，治療不應在這些部位上直接施行。
3. 可能需要局部麻醉。
4. 治療時，皮膚會有短暫的灼熱感及疼痛感覺。
5. 該療程所改善之程度及效果並沒有任何保證。效果因人而異，部份人可能未達到顯著之改善。
6. 若要得到較佳效果，需配合多次治療。
7. 治療前之部位需接受攝影，以作治療進度之參考。

手術 / 程序後須知

1. 治療後可繼續使用日常護膚產品（不適用於過紅及瘀傷之部位）
2. 治療後，部位會約有一星期的持續性輕微疼痛。

Patient's Label
Patient Name: _____
Hospital No:_____
Episode No:_____

風險及併發症

1. 治療之皮膚可能出現疼痛、刺痛、痕癢、泛紅、浮腫、鞭傷痕、瘀傷、灼傷、水泡、結疤、結焦、拉緊、疤痕形成、治療部位不對稱、神經受損及肌肉痲痺。
2. 治療之皮膚，色素可能變深或變淺，一般需時 1-3 個月或更長時間回復正常，情況因個人皮膚而異。
3. 當使用於減少脂肪時，可能會導致脂肪減少不均勻和皮膚表面不平整。

免責聲明

本單張只提供有關手術 / 程序的基本資料，並列出常見的風險或併發症；可能發生的風險或併發症不能盡錄，亦可因人而異。如需詳細資料或個別查詢，請聯絡你的醫生。

參考資料

1. Ulfit treatment guide.
2. Han HS et al. Safety and efficacy of high-intensity focused ultrasound for treatment of periorbital, perioral, and neck wrinkles: Prospective open single-center single-arm confirmatory clinical trial. Dermatologic Therapy. 2022;35:e15420.
3. Ayatollahi A et al. Systematic review and meta-analysis of safety and efficacy of high-intensity focused ultrasound (HIFU) for face and neck rejuvenation. Lasers in Medical Science. 2020;35:1007–1024.

病人簽署：_____ 簽署日期：_____